Frutex &c. Anguis &c.

ANGUIS GRACILIS CÆRULEO-VIRIDIS.

The blueiſh green Snake.

THIS harmleſs Snake frequents the Branches of Trees, and very nimbly ſquirms among the Leaves, living on Inſects, which I believe are its only Food. Their uſual Size and Length is that of the Figure. They are all over green, enclining more to blue than yellow. The Noſe of this Species turning up, ſufficiently diſtinguiſhes it from another green Snake I ſhall hereafter deſcribe.

Serpent d'un vert bleuâtre.

CE Serpent qui n'eſt point dangereux, ſe tient dans les branches des arbres, & ſe gliſſe fort legèrement entre les feuilles, vivant d'inſectes dont je croi qu'il fait ſa ſeule nourriture. Leur groſſeur & leur longueur ordinaire eſt ſemblable à la figure. Ils ſont par tout le corps d'un vert tirant plus ſur le bleu que ſur le jaune. Leur néz qui eſt tourné en haut, les diſtingue ſuffiſamment d'un autre ſerpent vert, dont je donnerai cy-après la deſcription.

Frutex baccifer, verticillatus; folijs ſcabris latis dentatis, & conjugatis; baccis purpurijs denſe congeſtis.

THESE Shrubs ariſe with ſeveral Stalks from the Ground, branching out on each Side, and in Height about four or five Feet; moſt of the Stalks are ſurrounded with Cluſters of very ſmall red Flowers, having four Petals each, with yellow *Stamina*, and growing at the Diſtance of about two Inches aſunder, beginning to flower at the Bottom, and ſucceſſively proceeding to flower to the Top of the Branch: From every Tuft of Flowers, grow oppoſite to each other a Pair of ferrated rough Leaves, the Berries which ſucceed the Flowers grow in Cluſters, ſo cloſely connected, that none of their Foot-ſtalks can be perceived without ſeparating them, which then diſcovers them to be held together by many ſmall branching Stalks. Theſe Berries are covered with a ſhining red Skin, containing many very ſmall Seeds. They bloſſom in *April* and *May*, the Berries are ripe in *July*. They grow in the Woods, near *Charles-Town* in *Carolina*.

CES arbriſſeaux s'elèvent de la terre par pluſieurs tiges, d'où ſortent des branches de chaque côté, ils croiſſent à la hauteur de quatre à cinq pieds: La pluſpart des tiges ſont entourées de bouquets de très petites fleurs rouges, ayant quatre feuilles avec des étamines jaunes, placés à la diſtance d'environ deux pouces les uns des autres. Ils commencent à fleurir par le bas, & continuent ſucceſſivement juſqu'au bout de la branche: De chaque bouquet de fleurs naiſſent à l'oppoſite l'une de l'autre deux feuilles rudes & dentelées. Les bayes qui ſuccèdent aux fleurs viennent en grapes; & ces bayes ſont ſi ſerrées les unes contre les autres, que ſans les ſeparer il eſt impoſſible d'appercevoir leurs pedicules; & alors vous les voyez attachées à de petits pedicules blancs. Ces bayes ſont couvertes d'une peau d'un rouge éclattant, & elles contiennent pluſieurs grains. Ces arbriſſeaux ſont en fleur au mois d'Avril & de Mai, *& leur fruit eſt mûr en* Juillet. *Ils croiſſent dans les bois, près de* Charles-Town *dans la* Caroline.